

REPORT DOCUMENTATION PAGE				<i>Form Approved OMB No. 0704-0188</i>							
The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.											
PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.											
1. REPORT DATE (DD-MM-YYYY) 2012		2. REPORT TYPE Journal Article-Experimental Physiology		3. DATES COVERED (From - To)							
4. TITLE AND SUBTITLE High skin temperature and hypohydration impair aerobic performance				5a. CONTRACT NUMBER							
				5b. GRANT NUMBER							
				5c. PROGRAM ELEMENT NUMBER							
6. AUTHOR(S) M.N. Sawka, S.N. Cheuvront, R.W. Kenefick				5d. PROJECT NUMBER							
				5e. TASK NUMBER							
				5f. WORK UNIT NUMBER							
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Thermal and Mountain Medicine Division U.S. Army Research Institute of Environmental Medicine Natick, MA 01760-5007				8. PERFORMING ORGANIZATION REPORT NUMBER MISC 11-47							
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Same as #7 above.				10. SPONSOR/MONITOR'S ACRONYM(S)							
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)							
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited.											
13. SUPPLEMENTARY NOTES											
14. ABSTRACT This paper reviews the roles of hot skin (>35°C) and body water deficits (>2% body mass; hypohydration) in impairing submaximal aerobic performance. Hot skin is associated with high skin blood flow requirements and hypohydration is associated with reduced cardiac filling, both of which act to reduce aerobic reserve. In euhydrated subjects, hot skin alone (with a modest core temperature elevation) impairs submaximal aerobic performance. Conversely, aerobic performance is sustained with core temperatures >40°C if skin temperatures are cool/warm when euhydrated. No study has demonstrated that high core temperature (-40°C) alone, without coexisting hot skin, will impair aerobic performance. In hypohydrated subjects, aerobic performance begins to be impaired when skin temperatures exceed 27°C, and even warmer skin exacerbates the aerobic performance impairment (-1.5% for each 1°C skin temperature). We conclude that hot skin (high skin blood flow requirements from narrow skin temperature to core temperature gradients), not high core temperature, is the 'primary' factor impairing aerobic exercise performance when euhydrated and that hypohydration exacerbates this effect.											
15. SUBJECT TERMS Hypohydration, submaximal aerobic performance, hypohydrated, euhydrated											
16. SECURITY CLASSIFICATION OF: <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 33%; padding: 2px;">a. REPORT</td> <td style="width: 33%; padding: 2px;">b. ABSTRACT</td> <td style="width: 33%; padding: 2px;">c. THIS PAGE</td> </tr> <tr> <td style="text-align: center;">Unclassified</td> <td style="text-align: center;">Unclassified</td> <td style="text-align: center;">Unclassified</td> </tr> </table>			a. REPORT	b. ABSTRACT	c. THIS PAGE	Unclassified	Unclassified	Unclassified	17. LIMITATION OF ABSTRACT Unclassified		18. NUMBER OF PAGES 6
a. REPORT	b. ABSTRACT	c. THIS PAGE									
Unclassified	Unclassified	Unclassified									
			19a. NAME OF RESPONSIBLE PERSON M.N. Sawka								
			19b. TELEPHONE NUMBER (Include area code) 508-233-5665								

Reset

Symposium Report

High skin temperature and hypohydration impair aerobic performance

Michael N. Sawka, Samuel N. Cheuvront and Robert W. Kenefick

Thermal and Mountain Medicine Division, US Army Research Institute of Environmental Medicine, Kansas Street, Natick, MA, USA

This paper reviews the roles of hot skin ($>35^{\circ}\text{C}$) and body water deficits ($>2\%$ body mass; hypohydration) in impairing submaximal aerobic performance. Hot skin is associated with high skin blood flow requirements and hypohydration is associated with reduced cardiac filling, both of which act to reduce aerobic reserve. In euhydrated subjects, hot skin alone (with a modest core temperature elevation) impairs submaximal aerobic performance. Conversely, aerobic performance is sustained with core temperatures $>40^{\circ}\text{C}$ if skin temperatures are cool-warm when euhydrated. No study has demonstrated that high core temperature ($\sim 40^{\circ}\text{C}$) alone, without coexisting hot skin, will impair aerobic performance. In hypohydrated subjects, aerobic performance begins to be impaired when skin temperatures exceed 27°C , and even warmer skin exacerbates the aerobic performance impairment (-1.5% for each 1°C skin temperature). We conclude that hot skin (high skin blood flow requirements from narrow skin temperature to core temperature gradients), not high core temperature, is the 'primary' factor impairing aerobic exercise performance when euhydrated and that hypohydration exacerbates this effect.

(Received 4 August 2011; accepted after revision 28 November 2011; first published online 5 December 2011)

Corresponding author R. W. Kenefick: Thermal and Mountain Medicine Division, US Army Research Institute of Environmental Medicine, 42 Kansas Street, Natick, MA 01760, USA. Email: robert.kenefick@us.army.mil

Heat stress impairs submaximal and maximal aerobic exercise performance (Sawka *et al.* 2011). For maximal intensity exercise, cardiovascular mechanisms related to oxygen delivery are likely to limit performance in the heat (Rowell *et al.* 1966; González-Alonso & Calbet, 2003). The mechanisms limiting sustained, submaximal intensity exercise in the heat include cardiovascular, CNS and metabolic (glycogen depletion) changes (Cheung & Sleivert, 2004; Sawka *et al.* 2011). Metabolic limitations are minor and specific to particular exercise tasks in the heat (Cheung & Sleivert, 2004). Cardiovascular mechanisms were historically assumed to be the primary factor impairing submaximal performance in the heat (Rowell, 1986), but the sustainment of skeletal muscle blood flow at exhaustion shifted the emphasis towards CNS limitations

and the role of high core temperatures (Nielsen *et al.* 1990, 1993).

Bodil Nielsen and colleagues (Nielsen *et al.* 1990) proposed that a high core temperature ($\sim 40^{\circ}\text{C}$) 'having an effect on the CNS in reducing the motor drive for performance' is the critical factor impairing submaximal intensity aerobic performance in the heat. During the past decade, the 'critical' core temperature hypothesis has been widely attributed as the primary mechanism impairing submaximal aerobic performance in the heat. Deteriorated CNS function may contribute to impaired aerobic performance in the heat (Nybo & Nielsen, 2001a), but the importance of a high core temperature has rarely been questioned. Studies supporting the 'critical' core temperature hypothesis have simultaneously induced high core temperatures with hot skin (Nielsen *et al.* 1990, 1993; González-Alonso *et al.* 1999). To our knowledge, no study has demonstrated that high core temperature alone will impair aerobic performance.

This paper reviews recent evidence that hot skin ($>35^{\circ}\text{C}$) alone can impair submaximal aerobic performance. Additional evidence will be provided

The opinions or assertions contained herein are the private views of the authors and are not to be construed as official or as reflecting the views of the Army or the Department of Defense. Any citations of commercial organizations and trade names in this report do not constitute an official Department of the Army endorsement of approval of the products or services of these organizations.

Table 1. Estimated whole-body skin blood flow (SkBF) requirements* during prolonged, severe running exercise† at different body core (T_c) and skin temperatures (T_{sk})

T_c (°C)	T_{sk} (°C)	Gradient (°C)	SkBF (l min ⁻¹)
38	30	8	1.1
38	32	6	1.5
38	34	4	2.2
38	36	2	4.4
39	30	9	1.0
39	32	7	1.3
39	34	5	1.8
39	36	3	2.9

*Equation for skin blood flow: $Q_s = 1/C \times h/(T_c - T_{sk})$, where C is the specific heat of blood ($\sim 0.87 \text{ kcal } ^\circ\text{C}^{-1} \text{ l}^{-1}$), h the heat production (in kcal min^{-1}) and Q_s the skin blood flow (Rowell, 1986). †Net heat production ($7.7 \text{ kcal min}^{-1}$) estimated using 60 kg body mass and 325 m min^{-1} running velocity (approximate pace for men's world class 42 km footrace) after subtracting for work (20% efficiency) and 50% dry and evaporative heat losses.

that if skin temperatures (T_{sk}) are cool-warm, aerobic performance can be sustained despite high core temperatures (T_c). Hot skin narrows the T_{sk} to T_c gradient, which increases skin blood flow requirements (Rowell, 1986) and may be the 'primary' factor impairing submaximal aerobic exercise performance in the heat. Body water deficits ($>2\%$ body mass; hypohydration) will exacerbate the effect by reducing central blood volume. We therefore postulate that during exercise heat stress, hot skin and hypohydration act in concert to reduce aerobic reserves, which increases the relative exercise intensity and perception of effort.

Physiology of skin temperatures and hypohydration

During exercise in the heat, the most significant physiological burden is to support high skin blood flow for heat dissipation (Sawka *et al.* 2011). Skin temperature is elevated in proportion to ambient temperature and humidity (Gagge & Gonzalez, 1996), while T_c is elevated in proportion to exercise intensity and is largely independent of the environment during compensable heat stress (Sawka *et al.* 2011). Warm-hot skin is associated with a greater skin blood flow and cutaneous venous compliance, which augments cardiovascular strain (Sawka *et al.* 2011). For this review, we define hot skin as 35°C and above, warm skin as $30\text{--}34.9^\circ\text{C}$ and cool/cold skin as $<30^\circ\text{C}$. We recognize that skin temperature effects are a continuum and that the T_{sk} to T_c gradient alters these relationships.

Table 1 illustrates the effects of different T_{sk} and T_c combinations on estimated (Rowell, 1986) whole-body skin blood flow requirements during combined exercise and heat stress. An elevated T_{sk} increases skin blood flow at any given T_c , while an elevated T_c reduces skin blood

flow requirements at any given T_{sk} . The rows beginning with T_c 38 and 39°C highlight an often unappreciated point; at any given skin temperature, an elevation in core temperature reduces whole-body skin blood flow and can be viewed as a positive response for sustaining aerobic performance in the heat. For example when comparing T_c of 39°C to a T_c of 38°C at equivalent T_{sk} of 36°C , SkBF is reduced from 4.4 to 2.9 l min^{-1} . Figure 1 demonstrates the impact of warm-hot T_{sk} , at constant T_c ($\sim 37.5^\circ\text{C}$), on cardiovascular strain during light-intensity (metabolic rate $\sim 450 \text{ W}$) exercise (Cheuvront *et al.* 2003). The heart rate (HR) elevation during exercise was an exponential function of skin warming beyond $T_{sk} \sim 35^\circ\text{C}$. The high skin blood flow requirements act to reduce cardiac filling and elevate HR for a given cardiac output (Trinity *et al.* 2010; Stohr *et al.* 2011b). Conversely, rapidly cooling T_{sk} has a profound effect on reducing HR and sustaining mean arterial pressure during exercise in the heat (Shaffrath & Adams, 1984). In addition, hot skin can be associated with reduced cerebral blood flow and cerebral oxygen delivery during moderate-intensity exercise (Nybo & Nielsen, 2001b; Nybo *et al.* 2002; Rasmussen *et al.* 2010).

During combined exercise and heat stress, hypohydration augments hyperthermia and cardiovascular strain in proportion to the magnitude of body water deficit (Sawka *et al.* 1985). Hypohydration reduces cardiac filling (Stohr *et al.* 2011b) and stroke volume during combined exercise and heat stress, making it difficult to maintain cardiac output (Montain & Coyle, 1992) and sustain muscle blood flow when heat stress is severe (González-Alonso *et al.* 1998).




Figure 1. Impact of high skin temperature, with a constant core temperature, on elevating heart rate during light-intensity exercise (metabolic rate $\sim 450 \text{ W}$)

From Cheuvront *et al.* (2003).

Aerobic performance

Heat stress. Warm-hot T_{sk} degrades maximal aerobic power ($\dot{V}_{O_{2max}}$) in proportion to the T_{sk} elevation (Arngrimsson *et al.* 2003). Thus, when performing exercise at a given metabolic rate, a person with warm-hot skin will work at a greater percentage $\dot{V}_{O_{2max}}$ compared with temperate conditions. Figure 2 demonstrates that marathon race performance is progressively slower with increased environmental (wet bulb globe temperature; WBGT) heat stress (Ely *et al.* 2007). Skin temperature is elevated with WBGT (Gagge & Gonzalez, 1996), but T_c may or may not be elevated, as it depends upon the sustainment of exercise intensity and heat exchange biophysics (Sawka *et al.* 2011). Therefore, marathon race performance might slow as a function of elevated T_{sk} .

Laboratory studies consistently demonstrate that T_{sk} elevations impair submaximal intensity aerobic performance. González-Alonso *et al.* (1999) employed a time-to-exhaustion (TTE; 60% $\dot{V}_{O_{2max}}$) test, during which subjects wore a water-perfused suit. When T_{sk} was elevated from ~ 36 to 38°C , TTE was shortened from 56 to 31 min; however, between the two trials, T_c ($\sim 40^\circ\text{C}$) and HR (~ 188 beats min^{-1}) were similar at exhaustion. MacDougall *et al.* (1974) used a similar combination of TTE test (70% $\dot{V}_{O_{2max}}$) with water-perfused suit to show that when T_{sk} was raised incrementally from ~ 29 to ~ 32 and then $\sim 35^\circ\text{C}$, TTE was shortened from 90 to 75 and then 48 min, respectively, despite similar core temperatures at exhaustion ($\sim 39.5^\circ\text{C}$). Tattersson *et al.* (2000) used a time-trial (TT) test and reported that performance was impaired by $\sim 6\%$ in a warm environment when T_{sk} was $\sim 33^\circ\text{C}$, versus 27°C in a temperate environment. Core temperature and HR levels were again similar at exhaustion ($\sim 39.3^\circ\text{C}$ and ~ 195 beats min^{-1}), as in other studies. Most recently, Periard *et al.* (2011) reported a $\sim 13\%$ decrement in mean power output during a 40 km TT in hot versus temperate environmental conditions that produced T_{sk} of 36 and 28°C , respectively. Although T_c was higher at exhaustion in the heat (39.8 versus 38.9°C), pacing strategy fell off significantly after 20 min of cycling when T_c was similar in both trials ($\sim 38^\circ\text{C}$), while T_{sk} was already $>5^\circ\text{C}$ higher in the heat.

Performance studies cited as directly supporting the ‘critical’ core temperature hypothesis have simultaneously elicited high core temperatures with hot skin. In the original study of Nielsen *et al.* (1993), subjects completed a TTE test (60% $\dot{V}_{O_{2max}}$) for 9–12 consecutive days in a hot environment as part of a heat acclimation experiment. Heat acclimation increased TTE from 48 to 80 min over the test days, with exhaustion consistently coinciding with $T_c \sim 40^\circ\text{C}$ and $T_{sk} \sim 37^\circ\text{C}$. González-Alonso *et al.* (1999) manipulated initial body temperatures prior to a TTE test (60% $\dot{V}_{O_{2max}}$) in a hot environment by applying

precooling, no precooling and preheating to subjects. The critical core temperature explanation for fatigue resulted from exhaustion coinciding with a consistently high T_c ($\sim 40^\circ\text{C}$), but T_{sk} ($\sim 37^\circ\text{C}$) and HR (~ 196 beats min^{-1}) were equally consistent, with the HR near maximal levels based on age.

There is evidence that hot T_{sk} ($>35^\circ\text{C}$) alone can degrade aerobic performance. Ely *et al.* (2010) measured the impact of two environmental conditions (40 and 20°C) on a 15 min TT performance test where T_c elevation was modest and similar in both trials ($\sim 38.2^\circ\text{C}$), but the compensable environments produced cool-warm (30°C) or hot skin (36°C). Time trial performance was impaired by 17% with hot T_{sk} , although a similar HR (~ 180 beats min^{-1}) was achieved. These findings are consistent with studies employing uncompensable heat stress to produce hot T_{sk} ($>35^\circ\text{C}$) while performing a walking TTE test (Sawka *et al.* 1992; Montain *et al.* 1994; Latzka *et al.* 1998). During those studies, physical exhaustion routinely occurred ($\sim 50\%$ of cases) at relatively low T_c ($<38.5^\circ\text{C}$), but with high HR relative to the exercise intensity. Therefore, hot skin will impair performance and induce exhaustion well below levels associated with the ‘critical’ T_c hypothesis.

There is evidence that competitive running performance (velocity) can be preserved despite high $T_c \geq 40^\circ\text{C}$, if T_{sk} is cool-warm (Ely *et al.* 2009; Lee *et al.* 2010). Ely *et al.* (2009) had highly trained runners perform an 8 km running TT on a 400 m track in compensable environmental conditions eliciting cool-warm T_{sk} (32 – 34°C). They measured running




Figure 2. Performance decrement (y-axis) based on marathon finishing time (x-axis) with increasing wet bulb globe temperature (WBGT)

This nomogram was developed by analysing weather data and finishers from seven marathons over many years (Ely *et al.* 2007).

velocities over 200 m segments and found no difference when T_c was below (first 6.5 km) or above 40°C (final 1.5 km). Figure 3 presents the individual data for average running velocities when T_c was below and above 40°C. Lee *et al.* (2010) also examined running performance in a warm environment during a longer, 21 km race, in which velocity was determined for 3 km intervals. They too found that high core temperature ($\geq 39.5^\circ\text{C}$) was common and not associated with reduced performance. Although T_{sk} was not measured, the T_{sk} prediction equation of Adams (1977) for outdoor running in the sun would suggest a value near 32°C.

Hypohydration. Hypohydration impairs maximal aerobic power in hot environments (Craig & Cummings, 1966) and submaximal aerobic performance in temperate and warm-hot environments (Cheuvront *et al.* 2005; Castellani *et al.* 2010; Kenefick *et al.* 2010). The following studies demonstrated that hypohydration impairs submaximal aerobic performance and that the impairment is augmented by high T_{sk} . In those studies, T_c was $<39^\circ\text{C}$ and therefore well below the 'critical' T_c .

Cheuvront *et al.* (2005) tested the effect of hypohydration on aerobic performance using a 30 min exercise preload at $\sim 50\% \dot{V}_{O_{2\max}}$, followed by a 30 min TT in temperate and cold environments. Hypohydration by 3% body mass impaired performance by 8% in the temperate ($T_{sk} \sim 29^\circ\text{C}$) but not in the cold environment ($T_{sk} \sim 20^\circ\text{C}$). Castellani *et al.* (2010) used a nearly identical test whereby T_{sk} was $\sim 32^\circ\text{C}$ in both hypohydration and euhydration trials. Hypohydration by 4% body mass impaired performance by 18%. Kenefick *et al.*




Figure 3. Time trial running velocities of 12 highly trained runners in compensable environmental conditions (cool-warm skin temperatures) when their core temperatures were below (mean of first ~ 32 200 m segments) or exceeded a core temperature of 40°C (mean of last approximately eight 200 m segments)

From Ely *et al.* (2009).

(2010) further characterized the interaction between environmental conditions and hypohydration by having subjects exercise for 30 min ($50\% \dot{V}_{O_{2\max}}$) followed by a 15 min TT in 10, 20, 30 and 40°C environments (inducing stepwise increases in T_{sk} from 26 to 36°C) when euhydrated and when hypohydrated by 4% body mass. Hypohydration impaired aerobic performance by 12 and 23% when T_{sk} was 33 and 36°C, respectively.

Figure 4 plots the impact of hypohydration on aerobic performance from the preceding three studies (Castellani *et al.* 2010; Cheuvront *et al.* 2005; Kenefick *et al.* 2010). These studies employed similar procedures over a broad range of T_{sk} from 20 to 36°C. Segmented regression (Vieth, 1989) was used to approximate the statistical T_{sk} threshold for performance impairment using individual study data points ($n = 53$ paired observations). The threshold which best minimized the residual sums of squares was shown to be 27.3°C, and warmer skin accentuated the performance impairment by $\sim 1.3\%$ for each additional 1°C rise in T_{sk} similar to that reported by Kenefick *et al.* (2010).

High skin temperature/relative intensity hypothesis

Cheuvront *et al.* (2010) proposed that impaired submaximal aerobic performance in the heat might be explained by warm-hot T_{sk} reducing $\dot{V}_{O_{2\max}}$. A large $\dot{V}_{O_{2\max}}$ is a prerequisite for success in sports where




Figure 4. Percentage decrement in submaximal aerobic performance from euhydration as a function of skin temperature (T_{sk}) when hypohydrated by 3–4% of body mass. Data are means (error bars are 95% confidence intervals) compiled from three studies (Cheuvront *et al.* 2005; Castellani *et al.* 2010; Kenefick *et al.* 2010) employing similar experimental procedures and time trial (TT) performance tests. Filled circles represent 15 min TT tests; open circles represent 30 min TT tests. At a T_{sk} intercept of $\sim 27^\circ\text{C}$, the percentage decrement in aerobic exercise performance declines linearly by $\sim 1.3\%$ for each 1°C rise in T_{sk} , similar to the single study of Kenefick *et al.* (2010). The best-fit equation for the second linear line segment is $y = -1.26x + 26.37$.

aerobic metabolism predominates (Bassett, & Howley, 2000), and $\dot{V}_{O_{2\max}}$ is reduced incrementally with warm-hot T_{sk} (Arngrimsson *et al.* 2003). When $\dot{V}_{O_{2\max}}$ is reduced, the resultant increased percentage $\dot{V}_{O_{2\max}}$ results in impaired submaximal exercise capacity (Gleser & Vogel, 1973a,b). If relative exercise intensity is increased, constant-rate exercise (TTE) will be more difficult to sustain (earlier fatigue) or require a slowing of self-paced exercise (TT) to achieve a similar sensation of effort. An increased percentage $\dot{V}_{O_{2\max}}$ is associated with greater cardiopulmonary stress (HR and respiration) and elevated perceived exertion, while warm-hot skin is associated with elevated thermal discomfort (Gagge *et al.* 1969; Gonzalez & Gagge, 1973). Other physiological cues being sensed might include cardiopulmonary baroreceptor unloading (Stohr *et al.* 2011a,b), reduced cerebral perfusion or cerebral oxygenation (Rasmussen *et al.* 2010) and arterial hypocapnia (Sawka *et al.* 1980). Likewise, if CNS function is deteriorated by heat stress and contributes to impaired performance, the afferent input from skin (Kunsch *et al.* 1995), muscle (Todd *et al.* 2005) and perhaps osmoreceptors/baroreceptors when hypohydrated (Montain & Tharion, 2010) might all contribute to altering the signal processing.

We conclude that: (1) hot skin (high skin blood flow requirements from narrow T_{sk} to T_c gradients), not high core temperature, is the 'primary' factor impairing submaximal aerobic performance when euhydrated; (2) hypohydration impairs submaximal aerobic performance when skin temperature is $\sim 27^\circ\text{C}$, and even warmer skin exacerbates (-1.5% for each 1°C T_{sk}) these decrements; and (3) high core temperature ($\sim 40^\circ\text{C}$) alone does not impair aerobic performance.

References

- Adams WC (1977). Influence of exercise mode and selected ambient conditions on skin temperature. *Ann NY Acad Sci* **301**, 110–127.
- Arngrimsson SA, Stewart DJ, Borrani F, Skinner KA & Cureton KJ (2003). Relation of heart rate to percent $\dot{V}_{O_{2\text{peak}}}$ during submaximal exercise in the heat. *J Appl Physiol* **94**, 1162–1168.
- Bassett DR Jr & Howley ET (2000). Limiting factors for maximum oxygen uptake and determinants of endurance performance. *Med Sci Sports Exerc* **32**, 70–84.
- Castellani JW, Muza SR, Cheuvront SN, Sils IV, Fulco CS, Kenefick RW, Beidleman BA & Sawka MN (2010). Effect of hypohydration and altitude exposure on aerobic exercise performance and acute mountain sickness. *J Appl Physiol* **109**, 1792–1800.
- Cheung SS & Sleivert GG (2004). Multiple triggers for hyperthermic fatigue and exhaustion. *Exerc Sport Sci Rev* **32**, 100–106.
- Cheuvront SN, Carter RI, Castellani JW & Sawka MN (2005). Hypohydration impairs endurance exercise performance in temperate but not cold air. *J Appl Physiol* **99**, 1972–1976.
- Cheuvront SN, Kenefick RW, Montain SJ & Sawka MN (2010). Mechanisms of aerobic performance impairment with heat stress and dehydration. *J Appl Physiol* **109**, 1989–1995.
- Cheuvront SN, Kolka MA, Cadarette BS, Montain SJ & Sawka MN (2003). Efficacy of intermittent, regional microclimate cooling. *J Appl Physiol* **94**, 1841–1848.
- Craig EN & Cummings EG (1966). Dehydration and muscular work. *J Appl Physiol* **21**, 670–674.
- Ely BR, Cheuvront SN, Kenefick RW & Sawka MN (2010). Aerobic performance is degraded, despite modest hyperthermia, in hot environments. *Med Sci Sports Exerc* **42**, 135–140.
- Ely BR, Ely MR, Cheuvront SN, Kenefick RW, DeGroot DW & Montain SJ (2009). Evidence against a 40°C core temperature threshold for fatigue in humans. *J Appl Physiol* **107**, 1519–1525.
- Ely MR, Cheuvront SN, Roberts WO & Montain SJ (2007). Impact of weather on marathon-running performance. *Med Sci Sports Exerc* **39**, 487–493.
- Gagge AP & Gonzalez RR (1996). Mechanisms of heat exchange: biophysics and physiology. In *Handbook of Physiology*, section 4, *Environmental Physiology*, ed. Fregly MJ & Blatteis CM, pp. 45–84. American Physiological Society, Bethesda, MD.
- Gagge AP, Stolwijk JA & Saltin B (1969). Comfort and thermal sensations and associated physiological responses during exercise at various ambient temperatures. *Environ Res* **2**, 209–229.
- Gleser MA & Vogel JA (1973a). Endurance capacity for prolonged exercise on the bicycle ergometer. *J Appl Physiol* **34**, 438–442.
- Gleser MA & Vogel JA (1973b). Effects of acute alterations of $\dot{V}_{O_{2\max}}$ on endurance capacity of men. *J Appl Physiol* **34**, 443–447.
- Gonzalez RR & Gagge AP (1973). Magnitude estimates of thermal discomfort during transients of humidity and operative temperature and their relation to the new ASHRAE effective temperature. *ASHRAE Trans* **79**, 88–96.
- González-Alonso J & Calbet JA (2003). Reductions in systemic and skeletal muscle blood flow and oxygen delivery limit maximal aerobic capacity in humans. *Circulation* **107**, 824–830.
- González-Alonso J, Calbet JA & Nielsen B (1998). Muscle blood flow is reduced with dehydration during prolonged exercise in humans. *J Physiol* **513**, 895–905.
- González-Alonso J, Teller C, Andersen SL, Jensen FB, Hyldig T & Nielsen B (1999). Influence of body temperature on the development of fatigue during prolonged exercise in the heat. *J Appl Physiol* **86**, 1032–1039.
- Kenefick RW, Cheuvront SN, Palombo LJ, Ely BR & Sawka MN (2010). Skin temperature modifies the impact of hypohydration on aerobic performance. *J Appl Physiol* **109**, 79–86.
- Kunsch E, Neccht S, Schnitzler A, Tyercha C, Schmitz F & Freund HJ (1995). Somatosensory evoked potentials elicited by intraneural microstimulation of afferent nerve fibers. *J Clin Neurophysiol* **12**, 476–487.
- Latzka WA, Sawka MN, Montain SJ, Skrinar GS, Fielding RA, Matott RP & Pandolf KB (1998). Hyperhydration: tolerance and cardiovascular effects during uncompensable exercise-heat stress. *J Appl Physiol* **84**, 1858–1864.

- Lee JK, Nio AQ, Lim CL, Teo EY & Byrne C (2010). Thermoregulation, pacing and fluid balance during mass participation distance running in a warm and humid environment. *Eur J Appl Physiol* **109**, 887–898.
- MacDougall JD, Reddan WG, Layton CR & Dempsey JA (1974). Effects of metabolic hyperthermia on performance during heavy prolonged exercise. *J Appl Physiol* **36**, 538–544.
- Montain SJ & Coyle EF (1992). Influence of graded dehydration on hyperthermia and cardiovascular drift during exercise. *J Appl Physiol* **73**, 1340–1350.
- Montain SJ, Sawka MN, Cadarette BS, Quigley MD & McKay JM (1994). Physiological tolerance to uncompensable heat stress: effects of exercise intensity, protective clothing, and climate. *J Appl Physiol* **77**, 216–222.
- Montain SJ & Tharion WJ (2010). Hypohydration and muscular fatigue of the thumb alter median nerve somatosensory evoked potentials. *Appl Physiol Nutr Metab* **35**, 456–463.
- Nielsen B, Hales JR, Strange S, Christensen NJ, Warberg J & Saltin B (1993). Human circulatory and thermoregulatory adaptations with heat acclimation and exercise in a hot, dry environment. *J Physiol* **460**, 467–485.
- Nielsen B, Savard G, Richter EA, Hargreaves M & Saltin B (1990). Muscle blood flow and muscle metabolism during exercise and heat stress. *J Appl Physiol* **69**, 1040–1046.
- Nybo L, Møller K, Volianitis S, Nielsen B & Secher NH (2002). Effects of hyperthermia on cerebral blood flow and metabolism during prolonged exercise in humans. *J Appl Physiol* **93**, 58–64.
- Nybo L & Nielsen B (2001a). Hyperthermia and central fatigue during prolonged exercise in humans. *J Appl Physiol* **91**, 1055–1060.
- Nybo L & Nielsen B (2001b). Middle cerebral artery blood velocity is reduced with hyperthermia during prolonged exercise in humans. *J Physiol* **534**, 279–286.
- Periard JD, Cramer MN, Chapman PG, Caillaud C & Thompson MW (2011). Cardiovascular strain impairs prolonged self-paced exercise in the heat. *Exp Physiol* **96**, 134–144.
- Rasmussen P, Nybo L, Volianitis S, Møller K, Secher NH & Gjedde A (2010). Cerebral oxygenation is reduced during hyperthermic exercise in humans. *Acta Physiol (Oxf)* **199**, 63–70.
- Rowell LB (1986). *Human Circulation: Regulation during Physical Stress*, Chapter 13: Circulatory adjustments to dynamic exercise and heat stress: competing controls. pp. 363–406. Oxford University Press, New York.
- Rowell LB, Marx HJ, Bruce RA, Conn RD & Kusumi F (1966). Reductions in cardiac output, central blood volume, and stroke volume with thermal stress in normal men during exercise. *J Clin Invest* **45**, 1801–1816.
- Sawka MN, Knowlton RG & Glaser RM (1980). Body temperature, respiration, and acid-base equilibrium during prolonged running. *Med Sci Sports Exerc* **12**, 370–374.
- Sawka MN, Leon LR, Montain SJ & Sonna LA (2011). Integrated physiological mechanisms of exercise performance, adaptation, and maladaptation to heat stress. *Compr Physiol* **1**, 1883–1928.
- Sawka MN, Young AJ, Francesconi RP, Muza SR & Pandolf KB (1985). Thermoregulatory and blood responses during exercise at graded hypohydration levels. *J Appl Physiol* **59**, 1394–1401.
- Sawka MN, Young AJ, Latzka WA, Neuffer PD, Quigley MD & Pandolf KB (1992). Human tolerance to heat strain during exercise: influence of hydration. *J Appl Physiol* **73**, 368–375.
- Shaffrath JD & Adams WC (1984). Effects of airflow and work load on cardiovascular drift and skin blood flow. *J Appl Physiol* **56**, 1411–1417.
- Stohr EJ, González-Alonso J, Pearson J, Low DA, Ali L, Barker H & Shave R (2011a). Effects of graded heat stress on global left ventricular function and twist mechanics at rest and during exercise in healthy humans. *Exp Physiol* **96**, 114–124.
- Stohr EJ, González-Alonso J, Pearson J, Low DA, Ali L, Barker H & Shave RE (2011b). Dehydration reduces left ventricular filling at rest and during exercise independent of twist mechanics. *J Appl Physiol* **111**, 891–897.
- Tattersson AJ, Hahn AG, Martin DT & Febbraio MA (2000). Effects of heat stress on physiological responses and exercise performance in elite cyclists. *J Sci Med Sport* **3**, 186–193.
- Todd G, Butler JE, Taylor JL & Gandevia SC (2005). Hyperthermia: a failure of the motor cortex and the muscle. *J Physiol* **563**, 621–631.
- Trinity JD, Pahnke MD, Lee JF & Coyle EF (2010). Interaction of hyperthermia and heart rate on stroke volume during prolonged exercise. *J Appl Physiol* **109**, 745–751.
- Vieth E (1989). Fitting piecewise linear regression functions to biological responses. *J Appl Physiol* **67**, 390–396.